

CLAIMS

What is claimed is:

- 5 1. An anti-microbial-adhesion pharmaceutical composition for administration to a patient in need of such a composition comprising an effective amount of an isolated adhesion inhibitory fraction from *Vaccinium* juice as an active ingredient and a pharmaceutically acceptable carrier or diluent, said isolated fraction having
- 10 (a) a molecular weight of $\geq 14,000$; and
 (b) anti-adhesion activity against *H. pylori*;
 (c) an elemental analysis of carbon 43-51%,
15 hydrogen 4 - 5%, no nitrogen, no sulfur and no chlorine;
 (d) an nuclear magnetic resonance (NMR) line spectrum as set forth in Figures 2A and 2B;
 (e) an ultraviolet spectrum with an absorption
20 peak at 280 nm in neutral or acidic pH solution which is absent in alkali solutions;
 (f) coaggregation reversal and coaggregation inhibition activity against oral bacteria and
 (g) an adhesion inhibitory activity against P
25 fimbriated bacteria; and
 wherein the concentration of the isolated adhesion inhibitory fraction is between 1 μ g and 10 mg per milliliter (ml).
- 30 2. A method of treating *H. pylori* infection in a patient in need of such treatment by administering to the patient an effective amount of the composition as set forth in claim 1.

3. A method of isolating a water extract fraction from a juice of berries of the *Vaccinium* plant genus exhibiting adhesion inhibitory activity against bacteria by the steps of

5 (a) dialyzing the juice extensively against double distilled water using dialysis tubing with a 14,000 molecular weight cut-off;

(b) lyophilizing the dialysate;

10 (c) fractionating the lyophilized dialysate on a polyacrylamide resin column; and

(d) eluting the fraction from the column with water.

15 4. The method of claim 3 wherein the fraction is eluted from the column with ammonia.

20 5. The method of claim 3 wherein the fraction eluted in step (d) is further purified by passage over a Fractogel column.

6. The method of claim 3 wherein the juice is cranberry juice.

25 7. The method of claim 6 wherein the cranberry juice is a commercially available preparation selected from the group consisting of juice and juice concentrate.

30 8. The method of claim 3 wherein the juice is blueberry juice.

9. An antibody directed against an isolated adhesion inhibitory water extract fraction from cranberry juice having

35 (a) a molecular weight of $\geq 14,000$;

(b) an elemental analysis of carbon 43 - 51%, hydrogen 4 - 5%, no nitrogen, no sulfur and no chlorine;

5 (c) a nuclear magnetic resonance (NMR) line spectrum as set forth in Figures 2A and 2B;

(d) an ultraviolet spectrum with an absorption peak at 280 nm in neutral or acidic pH solution which is absent in alkali solutions;

10 (e) an adhesion inhibitory activity against P fimbriated bacteria; and

(f) coaggregation reversal and coaggregation inhibition activity against oral bacteria.

15 10. An antibody as set forth in claim 9 selected from the group consisting of monoclonal and polyclonal antibody.

20 11. A fortified food composition providing anti-microbial-adhesion activity comprising a suitable food carrier and an effective amount of an isolated adhesion inhibitory water extract fraction from *Vaccinium* having

(a) a molecular weight of $\geq 14,000$;

(b) coaggregation reversal and coaggregation inhibition activity against oral bacteria

25 (c) an elemental analysis of carbon 43-51%, hydrogen 4 - 5%, no nitrogen, no sulfur and no chlorine;

(d) a nuclear magnetic resonance (NMR) line spectrum as set forth in Figures 2A and 2B;

30 (e) an ultraviolet spectrum with an absorption peak at 280 nm in neutral or acidic pH solution which is absent in alkali solutions; and

(f) an adhesion inhibitory activity against P fimbriated bacteria.

35

12. The composition as set forth in claim 11 wherein the food carrier is a fruit juice.

13. The composition as set forth in claim 12
5 wherein the fruit juice is cranberry juice.

14. An adhesion inhibitory fraction from a juice of berries of the *Vaccinium* plant genus exhibiting coaggregation reversal and coaggregation inhibition
10 activity against bacteria isolated by:

(a) dialyzing the juice extensively against double distilled water using dialysis tubing with a 14,000 molecular weight cut-off;

(b) lyophilizing the dialysate;

15 (c) fractionating the lyophilized dialysate on a polyacrylamide resin column; and

(d) eluting the fraction from the column with water.

20 15. The method of claim 14 wherein the fraction is eluted from the column with ammonia.

TE04240" 9849T660